

# F344 (Fischer 344)

## F344/NHsd

Developed in 1920 by Curtis and Dunning at the Columbia University Institute for Cancer Research, Irvington, NY, USA. The stock was purchased from a commercial breeder called "Fischer". In 1949 to Heston, National Cancer Institute, Bethesda, MD 20205, USA and in 1950 to Hansen, National Institutes of Health, Bethesda, MD 20205, USA. Subsequent sub-lines from either the Dunning or Hansen colonies.

## CHARACTERISTICS

The F344 rat is widely used in gerontology and in carcinogenicity testing programs.

### Animal model

The Fischer rat is useful for the following animal models: phenylketonuria (Anderson, 1982), interstitial cell tumor of the testis, urinary bladder carcinoma, esophageal carcinoma, and Lyme disease (Barthold *et al*, 1988).

### Anatomy

Unlike seven other strains it does not develop brownish skin scales on the dorsum of the body, the perineum and the tail (Tayama and Shisa, 1994). Low relative heart weight in ten weeks old males (Tanase *et al*, 1982).

## Behavior

Low open field defecation in males (Harrington, 1972). Low wheel activity (Harrington, 1971). Moderately easy to handle. Low response to operant morphine-reinforced behavior (Ambrosio *et al*, 1995). Low preference for ethanol and low capability to develop acute tolerance to ethanol hypnosis, like BN/Rij (York and Chan, 1994). Behavioral performance declined more rapidly with aging than in strain BN (Spangler *et al*, 1994). Develops smaller acoustic and tactile startle response than strain LEW, which may be associated with strain differences in hypothalamic-pituitary-adrenal activation (Glowa *et al*, 1992). Unusual in its lack of preference for any concentration of salt. However, there is a dramatic change from aversion to preference following transection of the chorda tympani nerve, which in other strains causes little change in salt preference (Sollars and Bernstein, 1994). A preference can be induced by depletion of sodium chloride (Breslin *et al*, 1995). F344 rats have been used in anxiety studies (Bert *et al*, 2001).

## Drugs

Neurological toxicity of polybrominated biphenyls and acrylamide described by Tilson *et al* (1978) and Tilson and Cabe (1979a,b). Low biliary excretion of copper after intravenous injection of CuSO<sub>4</sub> (Nederbragt and Lagerwerf, 1986). Susceptible to the induction of tumors of the tongue by 4-nitroquinoline-1-oxide

(Tanaka *et al*, 1993). Susceptible to the development of anaplastic astrocytomas and glioblastomas following treatment with N-methyl-N-nitrosourea in the drinking water (Shibutani *et al*, 1993).

Isolated tracheal rings hyperresponsive to carbachol compared with LEW and F344 (Jia *et al*, 1995). Relatively insensitive to the induction of tumors by N-methyl-N-nitrosourea (MNU) following treatment with cyproterone acetate, which caused a high incidence of tumors in the outbred Cpb:WU strain (Bosland *et al*, 1992a).

However, treatment with cyproterone acetate for 21 days, testosterone propionate for three days and a single i.p. injection of MNU results in atypical hyperplasia of the ventral prostate (Bosland *et al*, 1992b). Severity of spontaneous nephropathy in aged rats reduced by treatment with arylsulfonyleurea (Milman *et al*, 1979). Exposure of weanling rats to terephthalic acid or dimethyl terephthalate in the diet induced urolithiasis (Wolkowski *et al*, 1982). Phenylketonuria can be induced by p-Chloro-DL-phenylalanine with L-phenylalanine (Andersen and Guroff 1974; Anderson 1982). Bleomycin induces pulmonary fibrosis, which can be reduced by treatment with indomethacin (Thrall *et al*, 1979). Treatment with 3,2'-dimethyl-4-aminobiphenyl (DMAB) provides the best model for prostatic cancer of five tested (Shirai *et al*, 1990). Slow metabolizer of MPPB (Takahara *et al*, 1993). Morphine does not increase nose-poking behavior, and phenobarbital does not decrease it





(Witkin and Goldberg, 1992). Compared with LEW rats, F344 rats show a much lower preference for several classes of drugs of abuse. This may be associated with levels of neurofilament proteins in the ventral segmental area of the brain (Guitart *et al*, 1992). Duration of morphine-induced EEG slow-wave bursts and associated behavioral stupor was less in F344 than in LEW (Mayomichelson and Young, 1993a). F344 rats self-administer less morphine than LEW rats (Gosnell and Krahn, 1993). Duration of EEG slow-wave bursts and behavioral stupor also shorter in F344 than in LEW following administration of ethylketocyclazocine, suggesting differences in opioid-related receptor populations between these strains (Mayomichelson and Young, 1993b). Resistant to the organophosphate diisopropyl fluorophosphate (DFP) in terms of hypothermic response and recovery of day-night difference in core temperature (Gordon and Watkinson, 1995). Resistant (compared with Sprague Dawley rats) to the ventricular hypertrophy and pressure changes induced by monocrotaline. This is associated with pulmonary vascular response rather than hepatic metabolism (Pan *et al*, 1993). Oral administration of hydroquinone for two years resulted in dose-related nephropathy and renal tubule adenomas in males but not females, whereas Sprague Dawley rats were resistant (English *et al*, 1994). Sensitive to the convulsive effects of kainic acid (Golden *et al*, 1995). Relatively resistant to carcinogenic effects of tamoxifen, possibly associated with reduced cell proliferation in the liver (Carthew *et al*, 1995). Intermediate incidence of colorectal cancer after injections of azoxymethane (Kobaek-Larsen *et al*, 2002).

## Genetics

### Coat color genes

- *a, B, c, h* : albino.

### Histocompatibility

- *RT1<sup>M</sup>, RT2<sup>a</sup>, RT3<sup>b</sup>, RT8<sup>b</sup>*.

### Immunoglobulins

- *Igk-1<sup>b</sup>, Igh-1<sup>a</sup>, Igh-2<sup>a</sup>, Igh-3<sup>a</sup>* (Leslie, 1984).

### Biochemical markers

- *Acon-1<sup>b</sup>, Acp-2<sup>a</sup>, Ahd-2<sup>c</sup>, Akp-1<sup>a</sup>, Alb<sup>a</sup>, Amyl<sup>a</sup>, Cryg-1<sup>b</sup>, Es-1<sup>a</sup>, Es-2<sup>a</sup>, Es-3<sup>a</sup>, Es-4<sup>b</sup>, Es-6<sup>a</sup>, Es-7<sup>b</sup>, Es-8<sup>b</sup>, Es-9<sup>a</sup>, Es-10<sup>a</sup>, Es-14<sup>b</sup>, Es-15<sup>b</sup>, Es-16<sup>b</sup>, Es-18<sup>a</sup>, Fh<sup>b</sup>, Gc<sup>a</sup>, Glo-1<sup>a</sup>, Gox-1<sup>a</sup>, Hbb<sup>a</sup>, Igk-1<sup>b</sup>, Lap<sup>a</sup>, Mgd-1<sup>a</sup>, Mup-1<sup>b</sup>, Pep-3<sup>b</sup>, Pgd<sup>b</sup>*. (Bender *et al*, 1994).

## Immunology

Resistant to spontaneous autoimmune thyroiditis (Hajdu and Rona 1969), but susceptible to experimental allergic encephalomyelitis (Gasser *et al*, 1975), experimental allergic neuritis (Levine and Wenk, 1968), and autologous immune complex nephritis (Watson and Dixon, 1966). Relatively insensitive to the induction of experimental autoimmune glomerulonephritis (Sado *et al*, 1986). Susceptible to the development of experimental autoimmune myasthenia gravis (Biesecker and Koffler, 1988). Resistant to the induction of thyroiditis by 3-methylcholanthrene (Glover *et al*, 1969). Resistant to group A *Streptococcus pyogenes* and *Lactobacillus casei*-induced chronic polyarthritis (Lehman *et al*, 1984). Epitope specificities of collagen-induced arthritis studied by Cremer *et al*, (1992). Normally resistant to the development of adjuvant arthritis, but germ-free F344 rats are susceptible, and the use of paraffin oil rather than mineral oil also induces susceptibility

(Vandelangerijt *et al*, 1993). Low antibody response to streptococcal group A carbohydrate, not linked to the MHC (Stankus and Leslie, 1976). Neonatal pancreatic islets derived by non-enzymic (in vitro) isolation procedures can be transplanted across MHC barriers without any immune suppression, in contrast with other strains such as ACI (Ketchum *et al*, 1992). Tachykinins cause bronchoconstriction in susceptible F344 mainly by an indirect mechanism that involves stimulation of NK1 receptors and mast cell activation, in contrast with the less sensitive strain BDE where they cause bronchoconstriction by a direct effect on the airway smooth muscle via activation of NK2 receptors (Joos *et al*, 1994). Low primary and secondary response to sheep red blood cells (Tada *et al*, 1974).

Poor producers of reaginic antibody in response to ovalbumin in aluminium hydroxide (Murphy *et al*, 1974). Boulter and Sell (1984) have described the alphafetoprotein and albumin genes and compared these with ACI and BUF.

## Infection

Infection with *Hymenolepis diminuta* cysticercoids results in no worm loss and no mastocytosis in contrast with DA, where there was significant mastocytosis six weeks post infection and low persistence of worms (Ishih, 1992).

## Life-span and spontaneous disease

Life-span and tumor incidence depend both on strain characteristics and the environment. The following has been reported: Median lifespan about 31 months in males and 29 months in females with about 87% survival to 24 months in both sexes. (Sass *et al*, 1975).



Cameron *et al*, (1985) found a 75% survival at 26 months of age. Mean lifespan 24 months in both sexes in presence of severe pulmonary infection (Davey and Moloney, 1970). Median lifespan 23-31 months in barrier-reared males and 26-29 months in barrier-reared females (Sass *et al*, 1975, Coleman *et al*, 1977; Jacobs and Huseby, 1967; Hoffman, 1979; Yu *et al*, 1982). Food restriction to 60% of ad-libitum prolongs median lifespan to more than 34 months in males (Yu *et al*, 1982), but food restriction limited to early life and protein restriction caused only a small increase in longevity (Yu *et al*, 1985). Other studies of lifespan and neoplasia include Solleveld *et al* (1984) and Maekawa *et al* (1983). Most animals older than 2 years exhibit small local areas of nephritis; less than 25% show severe nephritis (Snell, 1967).

Tumors: Mammary tumors 41% in females and 23% in males, pituitary adenomas 36% in females and 24% in males, testicular interstitial cell tumors 85% in males. Other tumor types less common (Sass *et al*, 1975). Thyroid carcinoma 22% (Lindsey *et al*, 1968). Interstitial cell testicular tumors 65%, mononuclear cell leukemia 24%, subcutaneous fibroadenoma 9% in females. Both sexes have a 5% incidence of nodular hyperplasia of the liver. (Davey and Moloney, 1970). In various studies incidence of leukemia's 23-26% and of testicular interstitial tumors 65-90% (Jacobs and Huseby, 1967; Davey and Moloney, 1970; Moloney *et al*, 1970; Sass *et al*, 1975; Cockrell and Garer, 1976). Uterine polyploid tumors of endometrial origin 21% (Jacobs and Huseby, 1967). In germ-free conditions leukemia 26% in males, 36% in females, mammary tumors 12% in males 20% in females, all other tumors 9% in males, 5% in females (Sacksteder, 1976).

Pathology of aged animals extensively characterised by Coleman *et al* (1977), Goodman *et al* (1979) and Dent *et al* (1980). Aged rats exhibit peripheral retinal degeneration, which is exacerbated by fluorescent light of moderate intensity (32 foot-candles). They also develop cardiomyopathy with myocardial degeneration, fibrosis and chronic interstitial myocarditis (males 33%, females 18%) and nephropathy (67% in males, 39% in females) (Lai *et al*, 1979).

Retinas of both sexes show a steady decline with age in the thickness of the outer nuclear layer and photoreceptor layer, with a drastically accelerated rate of peripheral retinal degeneration seen only in males after 12 months of age (Diloreto *et al*, 1994; Faktorovich *et al*, 1992). Food restriction initiated at six months of age was as effective as food restriction initiated at 6 weeks of age in slowing the progression of chronic nephropathy and cardiomyopathy and in delaying the occurrence of neoplasia (Maeda *et al*, 1985).

Reduction of mineral or protein or fat intake without a reduction of energy had, at most, marginal effects on fatal neoplastic disease (Shimokawa *et al*, 1991).

### Miscellaneous

A detailed account of the pathology of F344 rats is given by Boorman *et al* (1990). Hematological parameters have been described by Lovell *et al* (1981). Effects of restraint, cage transportation, anesthesia and repeated bleeding on plasma glucose levels have been described by Tabata *et al* (1998). Characteristics of the COP strain have been described by Festing (1979) and Greenhouse *et al* (1990).

### Physiology and biochemistry

Growth described by Cameron *et al* (1985). Resistant to the development of salt-induced hypertension (Hall *et al*, 1976). High specific activity but low inducibility of NADP cytochrome C reductase compared with outbred Sprague Dawley rats (Gold and Widnell, 1975). Hepatic microsomal activity before and after induction by phenobarbitone well characterized (Page and Vesell, 1969; Gold and Widnell, 1975; Dent *et al*, 1980). Large pituitaries, susceptible to Cysticercus in infection and rapid absorption of diethylstilboestrol leading to death (Dunning *et al*, 1947). Low LD50 of pentobarbital sodium (70mg/kg) (Shearer *et al*, 1973). Have substantially higher levels of diurnal and stress-related corticosterone levels with higher levels of corticosteroid-binding globulin in plasma, spleen and thymus than LEW or Sprague Dawley rats (Dhabhar *et al*, 1993). Higher concentrations of cortical and hippocampal 5-HT1A receptors compared with LEW rats (Burnet *et al*, 1994). Hippocampal neurones are more vulnerable to ischemic insult than those of other strains (Iwasaki *et al*, 1995).

### Reproduction

Short gestation period:  $22.47 \pm .36$  days (Peters, 1986). Good breeding performance and large litter size. Males reach sexual maturity between 10 and 15 weeks, as determined by sperm production rate and other indicators of testicular and epididymal function (Blazak *et al*, 1985).



## REFERENCES

1. Ambrosio E, Goldberg SR, Elmer GI (1995) Behavior genetic investigation of the relationship between spontaneous locomotor-activity and the acquisition of morphine self-administration behavior. *Behavioral Pharmacology* 6, 229-237.
2. Anderson A, Guroff G (1974) Phenylketonuria. *Comp. Pathol. Bull.* 6, 1-2.
3. Anderson AE (1982) Phenylketonuria, supplemental update to model not 47. In: *Animal Models of Human Disease*, Fascicle 11 ed. (Capen CC, Hackel DB, Jones TC, Migaki G, eds) Registry of Comparative Pathology, Armed Forces Institute of Pathology, Washington, DC.
4. Barthold SW, Moody KD, Terwilliger GA, Jacoby RO, Steere AC (1988) An animal model for Lyme arthritis. *Ann. N.Y. Acad. Sci.* 539, 264-273.
5. Bender K, Balogh P, Bertrand MF, den Bieman M, von Deimling O, Eghtessadi S, Gutman GA, Hedrich HJ, Hunt SV, Kluge R, Matsumoto K, Moralejo DH, Nagel N, Portal A, Prokop C-M, Siebert RT, van Zutphen LFM (1994) Genetic characterization of inbred strains of the rat (*Rattus norvegicus*). *J. Exp. Anim. Sci.* 36, 151-165.
6. Bert B, Fink H, Sohr R, Rex A (2001) Different effects of diazepam in Fischer rats and two stocks of Wistar rats in tests of anxiety. *Pharmacology, Biochemistry and Behavior*, 70, 1-10.
7. Biesecker G, Koffler D (1988) Resistance to experimental autoimmune myasthenia gravis in genetically inbred rats. *J. Immunol.* 140, 3406-3410.
8. Blazak WF, Ernst LT, Stewart BE (1985) Potential indicators of reproductive toxicity: Testicular sperm production and epididymal sperm number, transit time, and motility in Fischer 344 rats. *Fund. Appl. Toxicol.* 5, 1097-1103.
9. Boorman GA, Eustis SL, Elwell MR, Montgomery JCA, MacKenzie WF (1990) Pathology of the Fischer rat. Academic Press, Inc, San Diego.
10. Bosland MC, Prinsen MK, Rivenson A, Silverman J, Fiala E, Williams GM, Kroes R, Weisburger JH (1992a) Induction of proliferative lesions of ventral prostate, seminal- vesicle, and other accessory sex glands in rats by N-methyl-N-nitrosourea - effect of castration, pretreatment with cyproterone- acetate and testosterone propionate, and rat strain. *Prostate* 20, 339-353.
11. Bosland MC, Prinsen MK, Rivenson A, Weisburger JH (1992b) Induction of skin and thyroid-tumors in male-rats by normal-methyl-normal-nitrosourea after sequential treatment with cyproterone-acetate and testosterone propionate - effects of castration, rat strain and time of carcinogen injection. *Carcinogenesis* 13, 669-674.
12. Boulter J, Sell S (1984) Polymorphism of rat alphafetoprotein and albumin genes. *Tumor Biol.* 5, 365-370.
13. Breslin PAS, Spector AC, Grill HJ (1995) Sodium specificity of salt appetite in Fischer-344 and Wistar rats is impaired by chorda tympani nerve transection. *American Journal of Physiology-Regulatory Integrative and Comparative Physiology* 38, R350-R356.
14. Burnet PWJ, Michelson D, Smith MA, Gold PW, Sternberg EM (1994) The effect of chronic imipramine administration on the densities of 5-HT<sub>1A</sub> and 5-HT<sub>2</sub> receptors and the abundances of 5-HT receptor and transporter messenger-RNA in the cortex, hippocampus and dorsal raphe of 3 strains of rat. *Brain Res.* 638, 311-324.
15. Cameron TP, Hickman RL, Kornreich MR, Tarone RE (1985) History, survival, and growth patterns of B6C3F<sub>1</sub> mice and F344 rats in the National Cancer Institute carcinogenesis testing program. *Fundam. Appl. Toxicol.* 5, 526-538.
16. Carthew P, Rich KJ, Martin EA, De Matteis F, Lim C-K, Manson MM, Festing MFW, White INH, Smith LL (1995) DNA damage as assessed by <sup>32</sup>P-postlabelling in three rat strains exposed to dietary tamoxifen: the relationship between cell proliferation and liver tumor formation. *Carcinogenesis* 16, 1299-1304.
17. Cockrell BY, Garner FM (1976) Interstitial cell tumor. *Comp. Pathol. Bull.* 8, 2-3.
18. Coleman GL, Barthold SW, Osbalsistan GW, Foster SJ, Jonas AM (1977) Pathological changes during aging in barrier-reared Fischer F344 male rats. *J. Gerontol.* 32, 258-278.
19. Cremer MA, Terato K, Watson WC, Griffiths MM, Townes AS, Kang AH (1992) Collagen-induced arthritis in rats - examination of the epitope specificities of circulating and cartilage-bound antibodies produced by outbred and inbred rats using cyanogen bromide-derived peptides purified from heterologous and homologous type-II collagens. *J. Immunol.* 149, 1045-1053.
20. Davey FR, Moloney WC (1970) Postmortem observations of Fischer rats with leukemia and other disorders. *Lab. Invest.* 23, 327-334.
21. Dent JG, Graichen ME, Schnell S, Lasker J (1980) Constitutive and induced hepatic microsomal cytochrome P-450 monooxygenase activities in male Fischer-344 rats and CD rats. A comparative study. *Toxicol. Appl. Pharmacol.* 52, 45-53.
22. Dhabhar FS, McEwen BS, Spencer RL (1993) Stress-response, adrenal-steroid receptor levels and corticosteroid-binding globulin levels - a comparison between Sprague-Dawley, Fischer-344 and Lewis rats. *Brain Res.* 616, 89-98.
23. DiIoreto D, Cox C, Grover DA, Lazar E, Delcerro C, Delcerro M (1994) The influences of age, retinal topography, and gender on retinal degeneration in the Fischer-344 rat. *Brain Res.* 647, 181-191.
24. Dunning WF, Curtis MR, Segaloff A (1947) Strain differences in response to diethylstilbestrol and the induction of mammary gland and bladder cancer in the rat. *Cancer Res.* 7, 511-521.
25. English JC, Perry LG, Vlaovic M, Moyer C, Odonoghue JL (1994) Measurement of cell-proliferation in the kidneys of Fischer-344 and Sprague-Dawley rats after gavage administration of hydroquinone. *Fundam. Appl. Toxicol.* 23, 397-406.
26. Faktovich EG, Steinberg RH, Yasumura D, Matthes MT, Lavail MM (1992) Basic fibroblast growth-factor and local injury protect photoreceptors from light damage in the rat. *Journal of Neuroscience* 12, 3554-3567.
27. Festing MFW (1979) Inbred strains. In: *The Laboratory Rat. Vol. I. Biology and Diseases* (Baker HJ, Lindsey JR, Weisbroth SH, eds). New York: Academic Press, pp 55-72.
28. Gasser DL, Palm J, Gonatas NK (1975) Genetic control of susceptibility to experimental allergic encephalomyelitis and the Ag-B locus of rats. *J. Immunol.* 115, 431-433.
29. Glover EL, Reuber MD, Godfrey F (1969) Methylcholanthrene-induced thyroiditis. Susceptibility of Buffalo strain rats. *Arch. Environ. Health* 18, 901-903.
30. Glowa JR, Geyer MA, Gold PW, Sternberg EM (1992) Differential startle amplitude and corticosterone response in rats. *Neuroendocrinology* 56, 719-723.
31. Gold G, Widnell CC (1975) Response of NADPH cytochrome c reductase and cytochrome P-450 in hepatic microsomes to treatment with phenobarbital-Differences in rat strains. *Biochem. Pharmacol.* 24, 2106.
32. Golden GT, Smith GG, Ferraro TN, Reyes PF (1995) Rat strain and age-differences in kainic acid-induced seizures. *Epilepsy Res.* 20, 151-159.
33. Goodman DC, Ward JM, Squire RA, Chu KC, Linhart MS (1979) Neoplastic and nonneoplastic lesions in aging F344 rats. *Toxicol. Appl. Pharmacol.* 48, 237-248.
34. Gordon CJ, Watkinson WP (1995) Strain differences in the laboratory rat - impact on the autonomic, behavioral, and biochemical response to cholinesterase inhibition. *J. Toxicol. Environ. Health* 45, 59-73.
35. Gosnell BA, Krahn DD (1993) Morphine-induced feeding - a comparison of the Lewis and Fischer 344 inbred rat strains. *Pharmacol. Biochem. Behav.* 44, 919-924.
36. Greenhouse DD, Festing MFW, Hasan S, Cohen AL (1990) Catalogue of inbred strains of rats. In: *Genetic monitoring of inbred strains of rats* (Hedrich HJ, ed). Stuttgart, New York: Gustav Fischer Verlag, pp. 410-480.
37. Guitart X, Beitnerjohnson D, Marby DW, Kosten TA, Nestler EJ (1992) Fischer and Lewis rat strains differ in basal levels of neurofilament proteins and their regulation by chronic morphine in the mesolimbic dopamine system. *Synapse* 12, 242-253.
38. Hajdu A, Rona G (1969) Spontaneous thyroiditis in laboratory rats. *Experientia* 25, 1325-1327.
39. Hall CE, Ayachi S, Hall O (1976) Immunity of Fischer 344 rats to salt hypertension. *Life Sci.* 18, 1001-1007.
40. Harrington GM (1971) Strain differences in rotating wheel activity of the rat. *Psychon. Sci.* 23, 363-364.
41. Harrington GM (1972) Strain differences in open-field behavior of the rat. *Psychon. Sci.* 27, 51-53.
42. Hoffman HJ (1979) Survival distributions for selected laboratory rat strains and stocks. In: *Development of the rodent as a model system for aging*. Book II ed. (Gibson DC, Adelman RC, Finch C, eds), pp. 19-34. DREW Pub. No. (NIH) 79-161, Washington, DC. (Department of Health, Education, and Welfare, Washington, D.C.: U.S.)
43. Ishii A (1992) Mucosal mast-cell response to *Hymenolepis diminuta* infection in different rat strains. *International Journal for Parasitology* 22, 1033-1035.
44. Iwasaki H, Ohmachi Y, Kume E, Kriegstein J (1995) Strain differences in vulnerability of hippocampal-neurons to transient cerebral-ischemia in the rat. *International Journal of Experimental Pathology* 76, 171-178.
45. Jacobs BB, Huseby RA (1967) Neoplasms occurring in aged Fischer rats, with special reference to testicular, uterine, and thyroid tumors. *J. Natl. Cancer Inst.* 39, 303-307.
46. Jia YL, Xu LJ, Heisler S, Martin JG (1995) Airways of a hyperresponsive rat strain show decreased relaxant responses to sodium-nitroprusside. *American Journal of Physiology-Lung Cellular and Molecular Physiology* 13, L85-L91.
47. Joos GF, Kips JC, Pauwels RA (1994) In-vivo characterization of the tachykinin receptors involved in the direct and indirect bronchoconstrictor effect of tachykinins in 2 inbred rat strains. *American Journal of Respiratory and Critical Care Medicine* 149, 1160-1166.
48. Ketchum RJ, Moore WV, Hegre OD (1992) Increased islet allograft survival after extended culture by a mechanism other than depletion of donor apcs - lack of correlation between the elimination of donor MHC class-II-positive accessory cells and increased transplantability. *Transplant.* 54, 347-351.
49. Kobaek-Larsen M, Fenger C, Hansen K, Nissen I, Diederichsen A, Thorup I, Den Bieman M, Vach W, Ritskes Holtinga M (2002) Comparative study of histopathologic characterization of azoxymethane-induced colon tumors in three inbred rat strains. *Comparative Medicine* 52, 50-57.
50. Lai Y-L, Jacoby RO, Yao PC (1979) Peripheral retinal degeneration in rats. *Am. J. Pathol.* 97, 449-452.
51. Lehman TJA, Allen JB, Plotz PH, Wilder RL (1984) Lactobacillus-casei cell wall-induced arthritis in rats. Cell-wall fragment distribution and persistence in chronic arthritis-susceptible LEW/N and arthritis-resistant F344/N rats. *Arthritis Rheum.* 27, 939-942.
52. Leslie GA (1984) Allotype determinants (Igh-3) associated with the IgG2c subclass of rat immunoglobulins. *Mol. Immunol.* 21, 577-580.
53. Levine S, Wenk EJ (1968) The production and transfer of allergic adrenitis. *Am. J. Pathol.* 52, 41-53.
54. Lindsey S, Nichols CW Jr, Chaikoff IL (1968) Naturally occurring thyroid carcinoma in the rat. Similarities to human medullary carcinoma. *Arch. Pathol.* 86, 353-364.
55. Lovell DP, Archer RK, Riley J, Morgan RK (1981) Variation in haematological parameters among inbred strains of rats. *Lab. Anim.* 15, 243-249.
56. Maeda H, Gleiser CA, Masoro EJ, Murata I, McMahan CA, Yu BP (1985) Nutritional influences on aging of Fischer 344 rats: II. Pathology. *J. Gerontol.* 40, 671-688.
57. Maekawa A, Kurokawa Y, Takahashi M, Kokubo T, Ogiu T, Odonera H, Tanigawa H, Ohono Y, Furikawa F, Hayashi Y (1983) Spontaneous tumors in F344/DuCr rats. *Gann* 74, 365-372.
58. Mayomichelson L, Young GA (1993a) EEG, EEG power spectral, and behavioral-differences in response to acute ethylketocyclazocine administration in 2 inbred rat strains. *Brain Res. Bull.* 31, 345-351.
59. Mayomichelson L, Young GA (1993b) Genetic profiles of morphine-induced EEG, EEG power spectra, and behavior in 2 inbred rat strains. *Brain Res.* 30, 79-84.
60. Milman HA, Peckham JC, Ward JM, Tyranny RE (1979) Reduction of the severity of nephropathy in aging Fischer 344 rats treated with analogs of arylsulfonyleurea. *Toxicol. Appl. Pharmacol.* 49, 425-430.
61. Moloney WC, Boschetti AE, King VP (1970) Spontaneous leukemia in Fischer rats. *Cancer Res.* 30, 41-43.
62. Murphy SM, Brown S, Miklos H, Fireman P (1974) Reagin synthesis in inbred strains of rats. *Immunol.* 27, 245-253.
63. Nederbragt H, Lagerwerf AJ (1986) Strain-related patterns of biliary excretion and hepatic distribution of copper in the rat. *Hepatology* 6, 601-607.
64. Page JG, Vesell ES (1969) Hepatic drug metabolism in ten strains of Norway rat before and after pre-treatment with phenobarbital. *Proc. Soc. Exp. Bio. Med.* 131, 256-261.
65. Pan LC, Wilson DW, Segall HJ (1993) Strain differences in the response of Fischer-344 and Sprague-Dawley rats to monocrotaline induced pulmonary vascular-disease. *Toxicology* 79, 21-35.
66. Peters AG (1986) Length of gestation period on eight inbred strains and three outbred strains of rats. *Animal Technology* 37, 109-112.
67. Sacksteder MR (1976) Occurrence of spontaneous tumors in the germfree F344 rat. *J. Natl. Cancer Inst.* 57, 1371-1373.
68. Sado Y, Naito M, Okigaha T (1986) Strain specific responses of inbred rats on the severity of experimental autoimmune glomerulonephritis. *J. Clin. Lab. Immunol.* 19, 193-199.
69. Sass B, Rabstein LS, Madison R, Nims RM, Peters RL, Kelloff GJ (1975) Incidence of spontaneous neoplasms in F344 rats throughout natural life-span. *J. Natl. Cancer Res.* 54, 1449-1456.
70. Shearer D, Creel D, Wilson CE (1973) Strain differences in the response of rats to repeated injections of pentobarbital sodium. *Lab. Anim. Sci.* 23, 662-664.
71. Shibutani M, Maekawa A, Okeda R, Mitsumori K, Imazawa T, Yoshida J, Onodera H, Hayashi Y (1993) An experimental-model for anaplastic astrocytomas and glioblastomas using adult F344 rats and N-methyl-N-nitrosourea. *Acta Pathologica Japonica* 43, 464-474.
72. Shimakawa L, Yu BP, Masoro EJ (1991) Influence of diet on fatal neoplastic disease in male Fischer 344 rats. *J. Gerontol.* 46, B228-B232.
73. Shirai T, Nakamura A, Fukushima S, Yamamoto A, Tada M, Ito N (1990) Different carcinogenic responses in a variety of organs, including the prostate, of five different rat strains given 3,2'-dimethyl-4-aminobiphenyl. *Carcinogenesis* 11, 793-797.
74. Snell KC (1967) Renal disease of the rat, in *Pathology of Laboratory Rats and Mice* (Cotchin E, Roe FJ, eds), pp. 105-147. Blackwell Scientific, Oxford.
75. Sollars SJ, Bernstein IL (1994) Gustatory deafferentation and desalination - effects on NaCl preference of Fischer-344 rats. *Am. J. Physiol.* 266, R510-R517.
76. Solleved HA, Haseman JK, McConnell EE (1984) Natural history of body weight gain, survival and neoplasia in the F344 rat. *J. Natl. Cancer Inst.* 72, 929-940.
77. Spangler EL, Waggle KS, Hengemihle J, Roberts D, Hess B, Ingram DK (1994) Behavioral assessment of aging in male Fischer 344 and brown Norway rat strains and their F1 hybrid. *Neurobiol. Aging* 15, 319-328.
78. Stankus RP, Leslie GA (1976) Rat interstrain antibody response and crossidiotypic specificity. *Immunogenet.* 3, 65-73.
79. Tabata H, Kitamura T, Nagamatsu N (1998) Comparison of restraint, cage transportation, anaesthesia and repeated bleeding on plasma glucose levels between mice and rats. *Lab. Anim.* 32, 142-148.
80. Tada N, Itakura K, Aizawa M (1974) Genetic control of the antibody response in inbred rats. *J. Immunogenet.* 1, 265-275.

81. Takahara E, Nagata O, Kato H, Ohta S, Hirobe M (1993) Interindividual differences of (+)-4-[4-(4-methylphenyl)phenylmethoxy-1-piperidinyl] butyric acid [(+)-mppb] disposition in rats. *Biol. Pharmaceut. Bull.* 16, 1057-1059.
82. Tanaka T, Kojima T, Kawamori T, Wang AJ, Suzui M, Okamoto K, Mori H (1993) Inhibition of 4-nitroquinoline 1-oxide-induced tongue carcinogenesis by the naturally-occurring plant phenolics caffeic, ellagic, chlorogenic and ferulic acids. *Carcinogenesis* 14, 1321-1325.
83. Tanase H, Yamori Y, Hansen CT, Lovenberg W (1982) Heart size in inbred strains of rats. Part 1. Genetic determination of the development of cardiovascular enlargement in rats. *Hypertension* 4, 864-872.
84. Tayama K, Shisa H (1994) Development of pigmented scales on rat skin - relation to age, sex, strain, and hormonal effect. *Lab. Anim. Sci.* 44, 240-244.
85. Thrall RS, McCormack JR, Jack RM, McReynolds RA, Ward PA (1979) Bleomycin-induced pulmonary fibrosis in the rat. Inhibition by indomethacin. *Am. J. Pathol.* 95, 117-130.
86. Tilson HA, Cabe PA (1979a) The effects of arylamide given acutely or in repeated doses on fore- and hindlimb function of rats. *Toxicol. Appl. Pharmacol.* 47, 253-260.
87. Tilson HA, Cabe PA (1979b) Studies on the neurobehavioral effects of polybrominated biphenyls in rats. *Ann. NY Acad. Sci.* 320, 325-336.
88. Tilson HA, Cabe PA, Mitchell CL (1978) Behavioral and neurological toxicity of polybrominated biphenyls in rats and mice. *Environ. Health Perspect.* 23, 257-263.
89. Vandellangerijt AGM, Vanlent PLEM, Hermus ARMM, Vandeputte LBA, Vandenberg WB (1993) Regulation of resistance against adjuvant arthritis in the Fischer rat. *Clin. Exp. Immunol.* 94, 150-155.
90. Watson JI, Dixon FJ (1966) Experimental glomerulonephritis. IX. Factors influencing the development of kidney in adjuvant nephritis in rats. *Proc. Soc. Exp. Biol. Med.* 121, 216-223.
91. Witkin JM, Goldberg SR (1992) Effects of D-amphetamine, Win-35,428, pentobarbital and morphine on schedule-controlled responding in 2 inbred rat strains that differ in locomotor stimulatory effects of cocaine. *Behavioral Pharmacology* 3, 455-463.
92. Wolkowski-Tyl R, Chin TY, Popp JA, Heck H d'A. (1982) Urolithiasis. *Am. J. Pathol.* 107, 419-421.
93. York JL, Chan AWK (1994) Absence of acute tolerance to ethanol hypnosis in F344 and BN/BiRij rats. *Alcohol* 11, 31-34.
94. Yu BP, Masoro EJ, McMahan CA (1985) Nutritional influences on aging of Fischer 344 rats. I: Physical, metabolic, and longevity characteristics. *J. Gerontol.* 40, 657-670.
95. Yu BP, Masoro EJ, Murata I, Bertrand HA, Lynd FT (1982) Life span study of Fischer 344 male rats fed ad libitum or restricted diets: Longevity, growth, lean body mass and disease. *J. Gerontol.* 37, 130-141.